


College of Arts and Sciences
Interdisciplinary Mathematics Institute
University of South Carolina

IMI Seminar Series



Comparing the Degrees of Unconstrained and Constrained Approximation By Polynomials

Dr. Dany Leviatan

Raymond and Beverly Sackler School of Mathematical Sciences, Tel Aviv University

It is quite obvious that one should expect that the degree of constrained approximation be worse than the degree of unconstrained approximation. However, it turns out that in certain cases we can deduce the behavior of the degrees of the former from information about the latter.

Let $E_n(f)$ denote the degree of approximation of $f \in C[-1, 1]$, by algebraic polynomials of degree $< n$, and assume that we know that for some $\alpha > 0$ and $N \geq 1$,

$$n^\alpha E_n(f) \leq 1, \quad n \geq N.$$

Suppose that $f \in C[-1, 1]$, changes its monotonicity or convexity $s \geq 0$ times in $[-1, 1]$ ($s = 0$ means that f is monotone or convex, respectively). We are interested in what may be said about its degree of approximation by polynomials of degree $< n$ that are comonotone or coconvex with f . Specifically, if f changes its monotonicity or convexity at $Y_s := \{y_1, \dots, y_s\}$ ($Y_0 = \emptyset$) and the degrees of comonotone and coconvex approximation are denoted by $E_n^{(q)}(f, Y_s)$, $q = 1, 2$, respectively. We investigate when can one say that

$$n^\alpha E_n^{(q)}(f, Y_s) \leq c(\alpha, s, N), \quad n \geq N^*,$$

for some N^* . Clearly, N^* , if it exists at all (we prove it always does), depends on α , s and N . However, it turns out that for certain values of α , s and N , N^* depends also on Y_s , and in some cases even on f itself.

<http://imi.cas.sc.edu/events/494/>

TUESDAY

April

22

4:00 - 5:00 PM

LeConte College

Room 312



UNIVERSITY OF
SOUTH CAROLINA
College of Arts and Sciences

The University of South Carolina is an equal opportunity institution.